

IOANNIS ANTONII SCOPOLI
MED. DOCT. S. C. R. ET APOST. MAIEST. MONTANÆ
CIVITATIS IDRÆ PHYSICI

ENTOMOLOGIA
CARNIOLICA
EXHIBENS
INSECTA
CARNIOLIÆ INDIGENA
ET
DISTRIBUTA IN ORDINES, GENERA,
SPECIES, VARIETATES.
METHODO LINNÆANA.



VINDOBONAE,
TYPIS IOANNIS THOMAE TRATTNER,
CAES. REG. AVLÆ TYPOGR. ET BIBLIOP.

MDCCLXIII.

In pratis paludotis, unibrosis, sylvestribus.

Oculi nigri. Antennarum articuli apicibus nigris. Corpus ferrugineum. Pedes pallide ferruginei. Huic similis in omnibus, sed alis uno puncto obsoleto, reperita circa aqueductus *Fodinarum Idriensis*, 14. Maii.

857. *TIPLA Vibratrix?*

— long. lin. 2½.

LINN. Faun. Svec. 2. 1761. quodammodo eadem.

Diagn. Corpus albo-virens, alæ albæ, oculi nigri, pedes motatorii.

Habitat circa paludes.

Culice non major. Alæ albæ immaculatæ. Pedes corpori concolores.

858. *TIPLA Fenestralis.*

— long. lin. 3.

Diagn. Alæ hyalinæ fusco-maculatæ. Abdomen fasciis fuscis (σ). Pedes ferruginei; geniculis fuscis.

In fenestris intra cubilia frequens, ex hortis involans.

Alæ costa, inter duas maculas fuscas marginales majores, flavescens. Oculi subvirides. Abdomen ferrugineum, fusco-fasciatum, villosulum.

859. *TIPLA Tremula.*

— long. lin. 3½.

LINN. Syst. Nat. p. 587.

Faun. Svec. 2. 1762. eadem descriptione.

Diagn. Alæ vivido-hyalinæ, immaculatæ. Thorax maculis (2-3), obscurioribus.

Pafsim occurrit.

Nostræ. Antennæ articulis æqualibus: primo crassiore. Thorax rufescens; maculis duabus, fuscis. Abdomen fuscum. Pedes colore thoracis. An ergo vere *Tremula LINNAEI*? Sit LINNÆUS ipse judex.