

The type material of Milichiidae and Carnidae (Insecta: Diptera: Schizophora) in the Naturhistorisches Museum Wien

I. Brake*

Abstract

The type specimens of Milichiidae and Carnidae in the Natural History Museum Vienna are listed. Lectotypes are designated for *Phylomyza epitacta* HENDEL, 1914, *Milichia integra* BECKER, 1922, and *Pholeomyia longifacies* HENDEL, 1933. Additionally, *Madiza palpalia* WAHLBERG is considered as an unavailable name for *Desmometopa discipalpis* PAPP, 1993.

Key words: Milichiidae, Carnidae, lectotype designation.

Zusammenfassung

Das Typenmaterial der Fliegenfamilien Milichiidae und Carnidae im Naturhistorischen Museum Wien wird aufgelistet. Lectotypen werden für die Arten *Phylomyza epitacta* HENDEL, 1914, *Milichia integra* BECKER, 1922 und *Pholeomyia longifacies* HENDEL, 1933 designiert. *Madiza palpalia* WAHLBERG wird als nicht verfügbarer Name für *Desmometopa discipalpis* PAPP, 1993 behandelt.

Introduction

The Diptera collection in the Naturhistorisches Museum Wien in Austria contains about 3 drawers of Milichiidae and 2 drawers of Carnidae. A visit to the museum in September 2005 revealed several unmarked types, most of which were described by Hendel.

The purpose of this paper is to list all Milichiidae and Carnidae types in the NHMW and to designate three lectotypes. The label data for all types are cited. Additionally, the species *Desmometopa discipalpis* PAPP, 1993 (synonym: *Madiza palpalia* WAHLBERG) is discussed, because the hitherto unknown female is present in NHMW. Photos of several Carnidae types can be found at <http://www.diptera.myspecies.info> and photos of several Milichiidae types can be found at <http://www.milichiidae.info>.

Terminology and classification of Milichiidae follow BRAKE (2000), the classification of Carnidae follows BRAKE (in press).

Abbreviations: H – handwritten, HT – holotype, LT – lectotype, P – printed, PLT – paralectotype, ST – syntype.

Type depositories: DEI – Deutsches Entomologisches Institut, Müncheberg, Germany;

* Dr. Irina Brake, Department of Entomology, The Natural History Museum, Cromwell Road, London SW7 5BD, United Kingdom – i.brake@nhm.ac.uk

HNHM - Hungarian Natural History Museum, Budapest, Hungary; NHMW - Naturhistorisches Museum Wien, Vienna, Austria; USNM – National Museum of Natural History, Washington, D.C., United States; UZMH - Zoological Museum, University of Helsinki, Finland.

Milichiidae

Phyllomyzinae

Aldrichiomyza agromyzina (HENDEL, 1911)

Aldrichiella agromyzina HENDEL, 1911: 37.

ST 3♂ 3♀ on one pin and 2♂ 3♀ on second pin. "Brookings / S Dak", "Aldrichia / agromyzina H. det. F. Hendel". Further ST are in the USNM.

Neophyllomyza leanderi (HENDEL, 1924)

Phyllomyza leanderi HENDEL, 1924: 406.

ST 2♂ 2♀ and two further ST covered in fungus. All specimens bear the label: "Austria sub. / Kremsm. / Czerny / [back of label:] 31/7.06". One male bears the additional label: "Phyllomyza / leanderi H. / ♂ det. Hendel" and one female bears the additional label: "Phyllomyza / leanderi H. / ♀ det. Hendel".

The type depository of two further syntypes mentioned in the original description is unknown.

Descriptive note: The halter of males is darker (nearly black) than the halter of females (nearly yellow).

Phyllomyza epitacta HENDEL, 1914

Phyllomyza epitacta HENDEL, 1914: 97.

ST 1♂, "Paroe, nördl. Paiwan- / Distr. (Formosa) / H. Sauter VIII. 1912", "Phyllomyza epitacta H. / Det. Hendel". There is an additional ♂ with the same locality label and a ♀ from "Chipun (Formosa) / H. Sauter VII. 1912".

Of these three specimens only the ♀ agrees with the description (yellow palpus, basoflagellomere yellow ventrally), which seems to be based on females only, as the enlarged basoflagellomere of males is not mentioned. According to HENNIG (1941) and ROHLFIEN & EWALD (1970) there was a syntype from Chipun in DEI, but the specimen is lost and only the pin remains. In contrast to the four above mentioned specimens/pins, Hendel's description is based on only 3 specimens from Chipun and Paroe. Based on the external evidence from Hendel's determination label I here designate the male from Paroe with Hendel's label as the lectotype. The purpose of this designation is to fix a male specimen as primary type because identification within this genus relies heavily on male characters. I cannot determine whether the female with the yellow palpus belongs to the same species.




Fig. 1: *Phyllomyza epitacta*, male epandrium with cercus and surstyli in lateral view and separately surstylus in greatest extension (posteroventral view). Abbreviations: app – appendix, ce – cercus, ep – epandrium, sur – surstylus. Scale bar: 0.1 mm.

Description of lectotype:

Coloration: Brown except for yellow distal tip of femora, fore- and midtibia, basal and distal tip of hindtibia, and all tarsi. Gena light brown, all setae reddish brown. Halter orangebrown, wing hyaline.

Head: Eye $1.4 \times$ as high as long, gena $0.1 \times$ as high as eye. Frons $1.3 \times$ as long as broad, frontal triangle extending to level of middle orbital seta. Chaetotaxy of frons as usual for *Phyllomyza*. Pedicel with one long seta, which is nearly as long as orbital setae, one seta which is less than half as long and several short setulae. Basoflagellomere $0.7 \times$ as long as eye, as long as high and nearly circular in lateral view except for slight apicoventral edge; covered with dense white pubescence about as long as base of arista thick. Arista inserted preapically on dorsal side, pubescence about $2 \times$ as long as base of arista thick. Palpus about as long as basoflagellomere, $3.4 \times$ as long as wide, flat, lanceolate, with short setulae at tip and ventral side, pubescence as long as on basoflagellomere, but brown and inconspicuous.

Thorax with brown microtomentum except for polished area on part of anepisternum and katepisternum; 1 postpronotal, 2 notopleural of which anterior seta is nearly $2x$ as long as posterior seta, 1 presutural, 1 supra-alar, 1 postalar, 1 intra-alar, and 3 postsutural dorsocentral setae, anterior dorsocentral only slightly longer than acrostichal setulae, acrostichal setulae in irregular rows, no prescutellar acrostichal seta, 1 basal and 1 cruciate apical scutellar seta, anepisternum bare, 1 katepisternal seta and row of setulae anterior to katepisternal seta from dorsal to ventral margin of katepisternum, no anepimeral seta.

Wing: 1.5 mm long, $2.3 \times$ as long as wide. Distal section of M₁ about $5 \times$ as long as penultimate section. Costal section between R₂₊₃ and R₄₊₅ nearly as long as section between R₄₊₅ and M₁.

Leg: Basitarsomere of hindleg with yellow comb of setulae.

Male terminalia (Fig. 1): Surstylus bilobate, distiphallus short, cercus with ventral appendix, which is directed posterad.

Phyllomyza epitacta is most similar to *P. nigripalpis* DE MEIJERE, but in this species the basoflagellomere apparently has no white pubescence and the halter is yellow. *Phyllomyza dilatata* MALLOCH, which also occurs in Taiwan, differs in having the hindfemur thickened and the palpus leaflike. *Phyllomyza epitacta* does not appear to have thickened hindfemora, but it is difficult to tell because the femora are flattened in both male specimens. The palpus is similar in both species, but according to the figure of Malloch it seems to be bigger in *P. dilatata*.

Phyllomyza equitans (HENDEL, 1919)

Neophyllomyza equitans HENDEL, 1919: 198.

ST 1♀, together with *Dolycoris baccarum* L. on one pin, "equitans / n.sp. / det. Hendel / [back of label:] 6./6. Bisamberg / W. Ö.", "Phyllomyza / equitans Hend. ♀ / Det. J. Swann 1997". ST 1♀, double mounted, same labels as first specimen but back side of first label reads: "Freiheitsau / Juli".

Phyllomyza longipalpis (SCHMITZ, 1924)

Neophyllomyza longipalpis SCHMITZ, 1924: 16.

HT ♂, "fenstr / Sittard / 20.8.19", "Neophyllomyza / longipalpis ♂ / Holotype Det. Schmitz", "longipalpis Schmitz".

Phyllomyza lucens HENDEL, 1924

Phyllomyza lucens HENDEL, 1924: 405.

HT ♀, "Rapallo. / 20/4. 4/302", "Phyll. / lucens H. / det. Hendel".

Phyllomyza melania (HENDEL, 1919)

Neophyllomyza melania HENDEL, 1919: 198.

HT ♀, "melania / n.sp. / det. Hendel / [back of label:] Unterberg / N. O. 16./6.". Additional 1♂ 3♀ from Austria, 1♂ 1♀ of these reared from "Eichenmulm" (rotting oak).

Phyllomyza tetragona HENDEL, 1924

Phyllomyza tetragona HENDEL, 1924: 407.

HT ♀, "Liegnitz / 4. 34045.", "Phyllomyza / tetragona / ♀ det. Hendel", "Phyllomyza tetragona Hendel ♀ / det. J. Swann 1997".

Madizinae

Desmometopa ciliata HENDEL, 1919

Desmometopa ciliata HENDEL, 1919: 200.

PLT 1♀, head missing, "Australia / Biró 1900", "N.S. Wales / Sydney", "Desmometopa / ciliata H. / det. Hendel", "Paratype [yellow label]".

The lectotype [HNHM] was designated by SABROSKY (1983).

Desmometopa singaporensis* KERTÉSZ, 1899Desmometopa singaporensis* KERTÉSZ, 1899: 194.

PLT 3♀, "Singapore, Biró 1898". One ♀ with additional label: "Desmometopa / singaporensis / Kert. [H] / det. Kertész" is a true *D. singaporensis*, but the ♀ with the additional label: "Desmometopa / singaporensis / Kert. det. Hendel" and the ♀ with the label: "singaporensis [H]" are in fact *D. varipalpis*.

These three specimens are probably the specimens mentioned by SABROSKY (1983: 45), even though they do not bear Sabrosky's label and even though one of them is a *D. singaporensis*, contrary to Sabrosky's statement that all three are *D. varipalpis*.

***Desmometopa sordida* (FALLÉN, 1820)**= *Agromyza m-atrum* MEIGEN, 1830: 170.

Possible ST 1♀, "atrum / Coll. Winth."

Desmometopa discipalpis* PAPP, 1993Desmometopa discipalpis* PAPP, 1993: 133.

1♂ 1♀ from Germany, Stuttgart, reared from "*Populus* Mulfm" (rotting poplar tree) with *Cossus*, determined by Hendel as *D. sordida*.

This is the first record of a ♀ and it differs from the male in having a normal shaped hindtibia. It does have enlarged palpus, forecoxa and -femur, but these are not as large as in the male. Contrary to PAPP's (1993: 134) statement, *D. discipalpis* keys to *D. sp. H* in SABROSKY's key (1983: 15), not *D. atypica* SABROSKY. The description of *Desmometopa* sp. H agrees with the description of *D. discipalpis* and Sabrosky's specimens were also reared "des galeries du *Cossus* de Moskat" in Algeria. Therefore I think that *Desmometopa* sp. H is *D. discipalpis*.

There is some confusion about the name *Madiza palpalia* WAHLBERG (in ZETTERSTEDT 1848: 2785). This name was introduced in a letter quoted by Zetterstedt in a footnote under *Agromyza sordida* FALLÉN. CHANDLER (1998: 143, CHANDLER et al. 2001: 201) used this name as a valid species-group name and as the possible senior synonym for *discipalpis* PAPP. Previously VERRALL in SCUDDER (1882: 215) recognized this name as a genus-group name, because it was written with a capital initial, as did SCHULZE & KÜKENTHAL (1935: 2468) and NEAVE (1939: 540). The name was not cited in the old Palearctic catalog by BECKER (1905), the new catalog by PAPP (1984), nor SABROSKY's (1983) revision of *Desmometopa*. However, BECKER (1907: 511) and HENNIG (1937: 43) used the name as a synonym to *Desmometopa sordida*. I consider this name to be unavailable according to ICZN Art. 11.6 (Publication as a synonym) because it was cited by Zetterstedt under *D. sordida* and because Wahlberg himself stated that *palpalia* is "perhaps" the male of *D. sordida*. As outlined above, the name has not been treated as an available name before 1961. Since the characters given by Wahlberg agree with the description of *Desmometopa discipalpis*, the valid name for this species is therefore *D. discipalpis*.

Leptometopa coquilletti* (HENDEL, 1907)Hypaspistomyia coquilletti* HENDEL, 1907: 241.

ST 1♀, "Aden, XII.98 / leg. O. Simony", "Hypsipistomyia / Coquillettii / Hendel". Also ST 1♀, "Aden, XII.98 / leg. O. Simony", "H. coquillettii / Hend. / det. Becker", "Hypsipistomyia H.".

Leptometopa lacteipennis (HENDEL, 1913)

Madiza lacteipennis HENDEL, 1913: 108.

ST 1♀, "Anping / Formosa / H. Sauter, V.1912", "Madiza ♀ / lacteipennis / H. det. Hendel". There is also a ♀ with the label: "V.12 Tainan / Formosa / H. Sauter", which could be a syntype.

In DEI there are two ST ♀ from Anping (ROHILFIEN & EWALD 1970). This would together make ST 3♀ from Anping. Hendel, however, cites 3♀ from Anping and Tainan, so the specimen from Tainan has to be a syntype and one specimen from Anping can't be a syntype.

Leptometopa latipes (MEIGEN, 1830)

Agromyza latipes MEIGEN, 1830: 177.

Possible ST 1♂, "Agromyza latipes M.". Also 1♀, "130", "[small green quadrangle]", "Coll. Winthem"; 1♂ "Hamb. / in fenestri", "Coll. Winthem"; 1♀, "Coll. Winthem"; 1♀, "[small green quadrangle]", "131", "Coll. Winthem".

Leptometopa rufifrons BECKER, 1903

= *Hypsipistomyia latigenis* HENDEL, 1933: 54.

HT ♂, "Hypsipist. / latigenis / ♂ Hend. / F. Hendel det. / [back of label:] Arbe / VII. 1928".

Milichiinae

Milichia decora (LOEW, 1870)

Lobioptera decora LOEW, 1870: 9.

6♂3♀ from Corsica in 1855 could be syntypes. Of these 4♂3♀ bear the labels: "Mann / Corsica / 1855 [P]", "decora Lw / det. F. Hendel", "speciosa Mg. / det. Schiner". 1♂ bears the label: "Mann / Cors. / 855 [H]" and 1♂ with abdomen missing the label: "decora Lw. / Corsica [H]".

Milichia integra BECKER, 1922

Milichia integra BECKER, 1922: 71.

ST 1♂, "Tonga / 10.-13.IV", "Aegypt. Sudan / Ebner, 1914", "M. integra / Beck. / det. Becker". ?ST 2♂1♀, "El Obeid / 13-14 III", "Aegypt. Sudan / Ebner, 1914".

The latter 3 specimens belong to another species, maybe *M. apicalis* SABROSKY. BECKER (1922) cites a type series of 1♂3♀ with the above data, so it is questionable, whether Becker made an error in citing the gender and the latter 3 specimens belong to the type series, or whether they do not belong. I here designate the specimen from Tonga as the lectotype in order to fix a specimen as primary type which corresponds to Becker's description.

Milichia speciosa* MEIGEN, 1830Milichia speciosa* MEIGEN, 1830: 132.

ST 1♂ 1♀, on one pin, "Marseille", "Coll. Winthem", "Milichia speciosa". ?ST 1♀, "speciosa", "Coll. Winthem".

= *Lobioptera marginata* MIK, 1864: 796.ST 1♀, "Illyria / Rubbia / 2.5.64 / Mik", "Lob. mar- / garitata mihi". Mik used both spellings, *marginata* and *margaritata* in his paper and BECKER (1905: 237) first revised the name and used *L. marginata* as the valid name.***Milichiella argyrogaster* (PERRIS, 1876)**= *Lobioptera tiefii* MIK, 1887: 178.

ST 3♂, "Carinthia / Ob. Federaun, Mik, 13.6.884", "Lobioptera tiefii Mik / det. Hendel". The male terminalia of one of the syntypes agree with the figures in IWASA (1999, fig. 7-9).

***Milichiella bimaculata* BECKER, 1907**= *Eccoptomma freyi* HENDEL, 1936: 6.

ST 1♀, "Tenerife / Icod. d. I. Vinos / R. Frey", "2251", "Eccoptomma / Freyi Hend.", "Typus?". This is one of three female syntypes described by HENDEL. FREY (1958: 62) incorrectly cited a holo- and allotype from UZMH, but these two specimens must be syntypes, too.

***Milichiella hendeli* BRAKE, 2000**= *Milichiella nitida* HENDEL, 1911: 39.

HT ♂, one wing missing, "Milichiella nitida n.sp. / det. F. Hendel", "Pacific Grove, Cal. / May 06", "wet meadow".

Pholeomyia anomala* HENDEL, 1933Pholeomyia anomala* HENDEL, 1933: 82.

HT ♂, in good condition, "13.3", "Soledade", "Bras. Exped. / Penther '03", "Pholeomyia anomala H. ♂ / F.Hendel det".

Pholeomyia argyrophenga* (SCHINER, 1868)Lobioptera argyrophenga* SCHINER, 1868: 291.

HT ♂, in bad condition, only thorax and wings left, "Lindig / 1864 / Venezuela", "Lobioptera / argyrophenga Schin.". This specimen was redescribed by HENDEL (1932: 135).

Pholeomyia leucozona* BILIMEK, 1867Pholeomyia leucozona* BILIMEK, 1867: 903.

LT ♂, dusted, wing tips damaged, "14. Jaenner", "Bilimek / Cacahua / milpa", "Pholeomyia leucozona", "LectoHOLOTYPE / Pholeomyia leucozona ♂ / 1958 C.W. Sabrosky [red label]".

Pholeomyia longifacies* HENDEL, 1933Pholeomyia longifacies* HENDEL, 1933: 81.

ST 1♂ 1♀, male in good condition, female slightly dusted. **The male is here designated as the lectotype** and has the following label data: "Brasilien", "Pholeomyia longifacies H. / F. Hendel det".

The purpose of this designation is to fix a male specimen as primary type because identification within this genus relies heavily on male characters. The female paralectotype has the following label data: "Fiebrig / Paraguay / S. Bernardino", "Pholeomyia longifacies H. / F. Hendel det".

Pholeomyia schineri* (HENDEL, 1932)Rhynchosmilichia schineri* HENDEL, 1932: 136.

ST 2♂, in good condition, "Brazil.", "argyrophenga / Schin. / det. Hendel", *Rhynchosmilichia / schineri* H. / F. Hendel det".

Carnidae***Carnus hemapterus* NITZSCH, 1818**

- = *Cenchridobia eggeri* SCHINER, 1862: 436.

- ST 1♂, "Austria / Coll. Egger. [P]", Eggeri [H] / det. Schiner [P]".

- = *Carnus setosus* STOBBE, 1913: 193.

- ST 2♂, "Zelebor / Picus / major / 1865 [H]".

Meoneura algerica* HENNIG, 1937Meoneura algerica* HENNIG, 1937: 61.

HT ♂, "Algeria centr., Hassi Babah".

Meoneura furcata* HENNIG, 1937Meoneura furcata* HENNIG, 1937: 64.

HT ♂, "Algeria centr., Hassi Babah".

***Meoneura obscurella* (FALLÉN, 1823)**

- = *Agromyza pectinata* MEIGEN, 1830: 179.

- ST 3♀, two in good condition, one worse, "pectinata [H] / Coll. Winthem"; ST 1?, abdomen and one wing missing, additional label "pectinata" in Meigen's handwriting. The type locality was not given by Meigen, it is possibly Austria.

- = *Agromyza infuscata* MEIGEN, 1830: 184.

- ST 1?, only head, thorax and rests of legs left, "infuscata [H] / Coll. Winthem", "infuscata [in Meigen's handwriting]". The type locality was not given by Meigen, it is possibly Austria.

Acknowledgements

I am very grateful to Peter Sehnal (NHMW) for his warm welcome and help. My thanks are due to the Schlinger Foundation for my former fellowship and the funding of the trip to Vienna. I thank Wayne N. Mathis and László Papp for reviewing the manuscript. I am very grateful to F. Christian Thompson for his help with the nomenclature of *Madiza palpalia* WAHLBERG.

References

- BECKER T., 1905: Cyclorrhapha Schizophora: Holometopa. – In: BECKER T., BEZZI M., KERTESZ K. & STEIN P. (eds.): Katalog der paläarktischen Dipteren. Vol. 4. – Budapest, pp. 1–273.
- BECKER T., 1907: Die Dipteren-Gruppe Milichiinae. – Annales historico-naturales Musei nationalis hungarici 5: 507–550.
- BECKER T., 1922: Wissenschaftliche Ergebnisse der mit Unterstützung der Akademie der Wissenschaften in Wien aus der Erbschaft Treitl von F. Werner unternommenen zoologischen Expedition nach dem anglo-ägyptischen Sudan (Kordofan) 1914. VI. Diptera. – Denkschriften der K. Akademie der Wissenschaften 98(1923): 57–82.
- BILIMEK D., 1867: Fauna der Grotte Cacahuamilpa in Mexiko. – Verhandlungen der Zoologisch-botanischen Gesellschaft in Wien 17: 901–908.
- BRAKE I., 2000: Phylogenetic systematics of the Milichiidae (Diptera, Schizophora). – Entomologica scandinavica, Supplements 57: 1–120.
- BRAKE I., in press: World catalog of Carnidae (Diptera, Schizophora). – Myia 12.
- CHANDLER P.J., 1998: Milichiidae. – In: CHANDLER P.J. (ed.): Diptera (incorporating a list of Irish Diptera). (Checklists of insects of the British Isles; 1). – Handbooks for the identification of British insects 12: 143–144.
- CHANDLER P.J., PETERSEN F.T. & PAPP L., 2001: Milichiidae. – In: PETERSEN F.T. & MEIER R. (eds.): A preliminary list of the Diptera of Denmark. – Steenstrupia 26: 201–202.
- FREY R., 1958: Kanarische Diptera brachycera p.p., von Håkan Lindberg gesammelt. – Commentationes biologicae 17(4): 1–63.
- HENDEL F., 1907: Neue und interessante Dipteren aus dem kaiserlichen Museum in Wien. – Wiener entomologische Zeitung 26: 223–245, 1 pl.
- HENDEL F., 1911: Über von Prof. J. M. Aldrich erhaltene und einige andere amerikanische Dipteren. – Wiener entomologische Zeitung 30: 19–46.
- HENDEL F., 1913: H. Sauter's Formosa-Ausbeute. Acalyptrate Musciden (Dipt.) II. – Supplementa entomologica 2: 77–112.
- HENDEL F., 1914: H. Sauter's Formosa-Ausbeute. Acalyptrate Musciden (Dipt.) III. – Supplementa entomologica 3: 90–117, 7 figs.
- HENDEL F., 1919: Neues über Milichiiden (Dipt.). – Entomologische Mitteilungen 8: 196–200.
- HENDEL F., 1924: Neue europäische *Phylomyza*-Arten (Dipt. Milichiidae). – Deutsche entomologische Zeitschrift 1924: 405–408.
- HENDEL F., 1932: Die Ausbeute der deutschen Chaco-Expedition. 1925/26. Diptera XXX–XXXVI. – Konowia. Zeitschrift für systematische Insektenkunde 11: 98–110, 115–145, 15 figs.
- HENDEL F., 1933: Neue acalyprate Musciden aus der paläarktischen Region (Dipt.). – Deutsche entomologische Zeitschrift 1933: 39–56.

- HENDEL F., 1936: [Agromyzidae & *E. Freyi* n. sp.]. – Die Dipterenfauna der Kanarischen Inseln und ihre Probleme: 120–124, 125–126.
- HENNIG W., 1937: 60a. Milichiidae et Carnidae. – In: LINDNER, E. (ed.): Die Fliegen der palaearktischen Region. – Schweizerbart, Stuttgart, Vol. 61a, pp. 1–91.
- HENNIG W., 1941: Verzeichnis der Dipteren von Formosa. – Entomologische Beihefte aus Berlin-Dahlem 8: iv+239 pp.
- IWASA M., 1999: The genus *Milichiella* Giglio-Tos (Diptera, Milichiidae) of Japan. – Medical entomology and zoology 50: 27–32.
- KERTÉSZ K., 1899: Verzeichnis einiger, von L. Biró in Neu-Guinea und am Malayischen Archipel gesammelten Dipteren. – Természetrajzi füzetek 22: 173–195.
- LOEW H., 1870: *Lobioptera speciosa* Meig. und *decora* nov. sp. – Zeitschrift für die gesamte Naturwissenschaft, N.F. 1 = 35: 9–14.
- MEIGEN J.W., 1830: Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten. – Schulz, Hamm, xi + 401 pp.
- MÍK J., 1864: Dipterologische Beiträge, I. Beschreibung neuer Arten. – Verhandlungen der Zoolisch-botanischen Gesellschaft in Wien 14: 791–797, Taf.21, Fig.5.
- MÍK J., 1887: Ueber Dipteren. I. Drei neue österreichische Dipteren. – Verhandlungen der Zoolisch-botanischen Gesellschaft in Wien 37: 173–180, Taf. IV.
- NEAVE S.A., 1939: Nomenclator zoologicus. A list of the names of genera and subgenera in zoology from the tenth edition of Linnaeus 1758 to the end of 1935. In four volumes. – Zoological Society of London, London, xiv + 957 pp.
- PAPP L., 1984: Family Milichiidae + Carnidae. – In: SOÓS Á. & PAPP L. (eds.): Clusiidae - Chloropidae. – Catalogue of Palaearctic Diptera 10, 110–123.
- PAPP L., 1993: Three new milichiid species from Hungary. – Annales historico-naturales Musei nationalis hungarici 85: 133–139.
- ROHLFIEN K. & EWALD B., 1970: Katalog der in den Sammlungen des ehemaligen Deutschen Entomologischen Institutes aufbewahrten Typen - VIII. – Beiträge zur Entomologie 22: 407–469.
- SABROSKY C.W., 1983: A Synopsis of the world species of *Desmometopa* Loew (Diptera, Milichiidae). – Contributions of the American Entomological Institute 19(8): 1–69.
- SCHINER J.R., 1862: Vorläufiger Commentar zum dipterologischen Theile der "Fauna austriaca". – Wiener entomologische Monatsschrift 6: 428–436.
- SCHINER J.R., 1868: Diptera (Art. 1). – In: WÜLLERSTORF-URBAIR B. (ed.): Reise der österreichischen Fregatte Novara 2 (Abt. 1, Sec. B). – Wien, 388 pp, 4 pls.
- SCHMITZ H., 1924: Een nieuwe vlieg uit Zuid-Limburg. – Natuurhistorisch maandblad 13: 16.
- SCHULZE F.E. & KÜKENTHAL W., 1926–40: Nomenclator animalium generum et subgenerum. – Preussische Akademie der Wissenschaften, Berlin, ccxlv + 3692 pp.
- SCUDDER S.H., 1882: Nomenclator zoologicus. Part 1. Supplemental list of genera in zoology. – Bulletin of the United States National Museum 19: xxi + 367 pp.
- STOBBE R., 1913: Zur Kenntnis der Gattung *Carnus* Nitzsch (*Cenchridobia* Schiner) mit 1 nov. sp. – Deutsche entomologische Zeitschrift 1913: 192–194.
- ZETTERSTEDT J.W., 1848: Diptera Scandinaviae disposita et descripta. Lundae, pp. 2581–2934.