



Far Eastern Entomologist

Дальневосточный энтомолог

Journal published by Far East Branch
of the Russian Entomological Society
and Laboratory of Entomology,
Institute of Biology and Soil Science,
Vladivostok

Number 279: 1-10

ISSN 1026-051X

June 2014

<http://urn:lsid:zoobank.org/pub:6321A35D-FD6E-4E0D-8AF9-9C1CD7360582>

NEW DATA ON THE SPIDER WASPS (HYMENOPTERA, POMPILIDAE) FROM THE WESTERN SIBERIA

R. T-o. Baghirov

*National research Tomsk state university Biological institute
Tomsk 634050, Russia. E-mail: rbaghirov@yandex.ru*

An annotated list of 36 species in 12 genera of the spider wasps collected in the Altay Republic and Altaysky krai in 2010–2012 is given. *Evagetes trispinosus* (Kohl, 1886) is newly recorded from Russia and 21 species are newly recorded from Western Siberia.

KEY WORDS: Hymenoptera, Pompilidae, spider wasps, distribution, Western Siberia.

Р. Т-о. Багиров. Новые данные о дорожных осах (Hymenoptera, Pompilidae) Западной Сибири // Дальневосточный энтомолог. 2014. № 279. С. 1-10.

Приведен аннотированный список 36 видов дорожных ос из 12 родов, собранных в Республике Алтай и Алтайском крае в 2010–2012 гг. *Evagetes trispinosus* (Kohl, 1886) указывается впервые для фауны России, а 21 вид – впервые для фауны Западной Сибири.

*Национальный исследовательский Томский государственный университет
Биологический институт, Томск, 634050, Россия.*

INTRODUCTION

The world fauna of Pompilidae numbers 4855 species in 125 genera and four subfamilies. There are 216 species in 32 genera and three subfamilies in Russia (Loktionov & Lelej, 2014). While the spider wasps fauna of some regions of the Europe, Central Asia, and Russia (European part and Far East) is studied rather well (Tobias, 1978; Wahis, 2013; Lelej & Loktionov, 2012; Loktionov & Lelej, 2014), the fauna of the Western Siberia is still not studied.

MATERIAL AND METHODS

The paper based on more than 80 specimens of spider wasps from Altay Republic and Altaysky krai deposited in the Zoological Institute (St. Petersburg), the Institute of Systematics and Ecology of Animals (Novosibirsk), and in the Tomsk State University (the author's collection).

The data for the places where the specimens were collected are as follows:

Chagan-Uzun – Altay Republic, Kosh-Agach area, Chagan-Uzun village, N 50°03'15" E 088°40'57" (Shcherbakov M.V.); **Chaynaya** – Altaysky krai, Tigireksky State Nature Reserve, the Chaynaya mountain (Baghirov R.T-o.); **Chemal** – Altay Republic, Chemalsky area, Chemal village, N 51°24'40" E 086°00'18" (Baghirov R.T-o.); **Kamlak** – Altay Republic, Shebalinsky area, Kamlak Gorno-Altaysk botanical Garden, N 51°37'28" E 085°42'10" (Baghirov R.T-o.); **Karakol** – Altay Republic, Onguday area, Karakol River pool, Karakol Nature Park "Uch Enmek", N 50°58'27" E 086°17'12" (Baghirov R.T-o.); **Kolyvansky** – Altaysky krai, Zmeinogorsky area, north slope of the Kolyvansky ridge, Kolyvanskoe Lake, N 51°21'50" E 082°11'30" (Baghirov R.T-o.); **Kozyr'** – Altaysky krai, Tigireksky State Nature Reserve, western exposure of the slope the Kozyr mountain (Baghirov R.T-o.); **Kuray** – Altay Republic, North Chuysky ridge, Kuray steppe, in the middle reaches Chuya River, N 50°12'39" E 087°54'18", (Shcherbakov M.V.); **Onguday** – Altay Republic, Onguday area, Onguday village, N 50°45'00" E 086°08'00" (Baghirov R.T-o.); **Tigirek** – Altaysky krai, Tigireksky State Natural Reserve, N 50°57'00" E 082°58'34" (Baghirov R.T-o.).

New distributional data are asterisked (*).

LIST OF THE SPECIES

Subfamily Ceropalinae Radoszlowski, 1888

Ceropales (Ceropales) helvetica Tournier, 1889

MATERIAL. Onguday: 25.VII 2012, 1 ♀; Chemal: 28.VII 2012, 1 ♀.

DISTRIBUTION. Russia (Transcaucasia, south, south-east of European part, *Altay Republic), Europe, Central Asia (Wahis, 2013; Tobias, 1978).

Subfamily Pepsinae Lepeletier de Saint Fargeau, 1845

Auplopus carbonarius (Scopoli, 1763)

Auplopus carbonarius: Lelej, 1995: 233 (key), ♀ ♂.

MATERIAL. Kozyr': 11.VII 2012, 1 ♀.

DISTRIBUTION. Russia (European part, *Altaysky krai, Irkutskaya oblast, Primorskii krai, Amurskaya oblast, Khabarovskii krai, Magadanskaya oblast), Western Europe, Ukraine, Kazakhstan, Near East, Korea, Japan, North Africa (Lelej & Loktionov, 2012).

Caliadurgus fasciatellus (Spinola, 1808)

MATERIAL. Kamlak: 18.VII 2012, 1 ♀, 1 ♂; Chaynaya: 09.VII 2012, 1 ♀.

DISTRIBUTION. Russia (European part, *Altay Republic, Irkutskaya oblast, Buryatia, Sakhalin, Primorskii krai, Amurskaya oblast, Khabarovskii krai, Kamchatka), Western Europe, Ukraine, Belarus (Lelej, Loktionov, 2012).

Cryptocheilus (Adonta) fabricii (Vander Linden, 1827)

MATERIAL. Kozyr': 11.VII 2012, 1 ♀.

DISTRIBUTION. Russia (Caucasus, European part, *Altaysky krai, Irkutskaya oblast), Western Europe, Ukraine, Kazakhstan (Tobias, 1978; Lelej, 1995; Wahis, 2013).

Cryptocheilus (Adonta) freygessneri (Kohl, 1883)

MATERIAL. Chemal: 28.VII 2012, 1 ♀.

DISTRIBUTION. Russia (south and south-east of European part, *Altay Republic), Europe (Czech Republic, Italian mainland, Romania, Slovakia) (Tobias, 1978; Wahis, 2013).

Deuteragenia bifasciata (Geoffroy, 1785)

Dipogon hircanum: Tobias, 1978: 92 (key), ♀ ♂.

Dipogon bifasciatus: Lelej, 1995: 228 (key), ♀ ♂.

Deuteragenia bifasciata: Lelej & Loktionov, 2012: 413.

MATERIAL. Chemal: 28.VII 2012, 1 ♀; Kozyr': 11.VII 2012, 1 ♂.

DISTRIBUTION. Russia (European part, West Siberia, Irkutskaya oblast, Buryatia, Zabaikalskii krai, Sakhalin, Primorskii krai, Amurskaya oblast, Khabarovskii krai, Magadanskaya oblast), Western Europe, Belarus, Ukraine, Central Asia, Korea, Japan (Hokkaido, Honshu, Kyushu) (Lelej & Loktionov, 2012).

Deuteragenia vechti (Day, 1979)

Dipogon bifasciatum Geoffroy: Tobias, 1978: 91 (key), ♀ ♂.

Dipogon vechti Day: Lelej, 1995: 228 (key), ♀ ♂.

Deuteragenia vechti (Day, 1979): Lelej & Loktionov, 2012: 413.

MATERIAL. Chaynaya: 09.VII 2012, 2 ♀; Kamlak 18.VII 2012, 1 ♀.

DISTRIBUTION. Russia (European part, Siberia, Irkutskaya oblast, Khabarovskii krai, Amurskaya oblast, Jewish autonomous oblast, Primorskii krai, Sakhalin, South Kuril Islands, Kamchatka, Magadanskaya oblast, Chukotka), Western Europe, Belarus, Central Asia, Japan (Hokkaido, Honshu) (Lelej & Loktionov, 2012).

***Priocnemis (Priocnemis) agilis* (Shuckard, 1837)**

Priocnemis (P.) obtusiventris: Tobias, 1978: 104, ♀ ♂.

MATERIAL. Tigirek: 05.VIII 2011, 1 ♀; Kolyvansky: 06.VII 2011, 1 ♀.

DISTRIBUTION. Russia (Transcaucasia, European part, *Altaysky krai), Western Europe, Central Asia (Tobias, 1978; Wahis, 2013).

***Priocnemis (Priocnemis) exaltata* (Fabricius 1775)**

Priocnemis (P.) exaltata: Tobias, 1978: 102 (key), ♀ ♂.

MATERIAL. Tigirek: 05.VIII 2011, 1 ♀.

DISTRIBUTION. Russia (European part, *Altaysky krai), Western Europe, Ukraine (Tobias, 1978; Wahis, 2013).

***Priocnemis (Priocnemis) fenestrata* (Gussakovskij, 1926)**

MATERIAL. Kamlak: 18.VII 2012, 1 ♀; Chemal: 28.VII.2012, 1 ♀.

DISTRIBUTION. Russia (*Altay Republic, Irkutskaya oblast, South Yakutia, Primorskii krai, Amurskaya oblast, Khabarovskii krai), Republic of Korea (Lelej & Loktionov, 2012).

***Priocnemis (Priocnemis) hyalinata* (Fabricius, 1793)**

Priocnemis (P.) femoralis: Tobias, 1978: 100, ♀ ♂.

MATERIAL. Kamlak: 18.VII 2012, 1 ♀.

DISTRIBUTION. Russia (European part, *Altay Republic), Western Europe, Uzbekistan (Tobias, 1978; Wahis, 2013).

Subfamily Pompilinae Latreille, 1804

***Anoplius (Anoplius) aberrans* Gussakovskij, 1932**

MATERIAL. Kozyr': 11.VII 2012, 1 ♀.

DISTRIBUTION. Russia (*Altaysky krai, Irkutskaya oblast, Buryatia, Kuril Islands, Sakhalin, Primorskii krai, Amurskaya oblast, Jewish autonomous oblast, Khabarovskii krai, Magadanskaya oblast, Kamchatka), Uzbekistan, Japan (Hokkaido, Honshu), Republic of Korea, China (Guangdong, Taiwan), Philippine (Luzon), Indonesia (Celebes) (Loktionov & Lelej, 2012).

Anoplius (Anoplius) concinnus (Dahlbom, 1843)

MATERIAL. Chermal: 28.VII 2012, 1 ♀.

DISTRIBUTION. Russia (northwest and south of European part, *Altay Republic, Irkutskaya oblast, Buryatia, Primorskii krai, Amurskaya oblast, Khabarovskii krai), Europe, Kazakhstan, Central Asia, Iran, Mongolia (Shlyahtenok *et al.*, 2012).

Anoplius (Anoplius) nigerrimus (Scopoli, 1763)

MATERIAL. Kozyr': 11.VII 2012, 1 ♂.

DISTRIBUTION. Russia (European part, *Altaysky krai, Irkutskaya oblast, Buryatia, South Yakutia, Sakhalin, Kuril Islands, Primorskii krai, Amurskaya oblast, Khabarovskii krai, Kamchatka), Europe, North America (Shlyahtenok *et al.* 2012).

Anoplius (Anoplius) tenuicornis (Tournier, 1889)

Anoplius (Anoplius) tenuicornis: Tobias, 1978: 120 (key), ♀ ♂ (East Siberia).

MATERIAL. Chermal: 28.VII 2012, 2 ♀.

DISTRIBUTION. Russia (European part, *Altay Republic, Krasnoyarskii krai, Irkutskaya oblast, Sakhalin, Primorskii krai, Khabarovskii krai, Magadanskaya oblast, Kamchatka), Europe, Central Asia, Mongolia, North America (Shlyahtenok *et al.*, 2012).

Anoplius (Arachnophroctonus) viaticus (Linnaeus, 1758)

Anoplius (P.) viaticus: Tobias, 1978: 120 (key), ♀ ♂; Baghirov, 2011: 219, ♀.

MATERIAL. Kozyr': 11.VII 2012, 1 ♂; Chaynaya: 09.VII 2012, 3 ♂; Tigirek: 05.VIII 2011, 1 ♀, 6 ♂.

DISTRIBUTION. Russia (European part, Altaysky krai, Irkutskaya oblast, Buryatia, South Yakutia, Zabaikalskii krai, Sakhalin, Primorskii krai, Amurskaya oblast, Jewish autonomous oblast, Khabarovskii krai), Europe, Kazakhstan, Turkmenistan, Mongolia, Japan (Honshu, Kyushu), South Korea, Georgia, North Africa (Shlyahtenok *et al.*, 2012).

Aporinellus sexmaculatus (Spinola, 1805)

Aporinellus sexmaculatus: Tobias, 1978: 122 (key), ♀ ♂.

MATERIAL. Chermal: 28.VII 2012, 1 ♀, 1 ♂.

DISTRIBUTION. Russia (Caucasus, European part, south-west Siberia, Altay Republic), Europe, Near East, North Africa (Tobias, 1978; Wahis, 2013).

Arachnospila (Ammosphex) anceps (Wesmael, 1851)

MATERIAL. Kamlak: 18.VII 2012, 1 ♂.

DISTRIBUTION: Russia (European part, *Altay Republic, Buryatia, Zabaikalskiy krai, Amurskaya oblast, Chukotka, Magadanskaya oblast, Kamchatka), Europe, Turkey, Syria, Iran, Kyrgyzstan, Mongolia (Loktionov & Lelej, 2011).

***Arachnospila (Ammosphex) kurzenkoi* Lelej, 1995**

MATERIAL. Kozyr': 11.VII 2012, 1 ♂.

DISTRIBUTION. Russia (*Altaysky krai, Buryatia, Primorskiy krai, Amurskaya oblast, Khabarovskii krai, Chukotka, Magadanskaya oblast) (Loktionov & Lelej, 2011).

***Arachnospila (Ammosphex) mongolopinata* Wolf, 1981**

MATERIAL. Kozyr': 11.VII 2012, 2 ♂.

DISTRIBUTION. Russia (*Altaysky krai, Sakhalin, Primorskiy krai, Amurskaya oblast), Mongolia (Mongolian and Gobi Altay) (Loktionov & Lelej, 2011).

***Arachnospila (Ammosphex) trivialis* (Dahlbom, 1843)**

Arachnospila (Ammosphex) trivialis: Tobias, 1978: 128 (key), ♀ ♂; Lelej, 1995: 246 (key), ♀ ♂; Loktionov & Lelej, 2011: 24, ♀ ♂.

MATERIAL. Kozyr': 11.VII 2012, 1 ♂.

DISTRIBUTION. Russia (European part, Siberia, Chukotka, Magadan oblast, Kamchatka), Europe, Kazakhstan, Kyrgyzstan (Loktionov & Lelej, 2011).

***Arachnospila (Ammosphex) minutula* (Dahlbom, 1842)**

Arachnospila (Ammosphex) minutula: Tobias, 1978: 132 (key), ♀ ♂.

MATERIAL. Kozyr': 11.VII 2012, 1 ♀.

DISTRIBUTION. Russia (European part, Siberia, Buryatia, Zabaikalskii krai, Primorskii krai, Amurskaya oblast, Khabarovskii krai), Western Europe, Ukraine, Belarus, Mongolia, Near East (Lelej & Loktionov, 2012).

***Arachnospila (Anoplochares) mongolospissa* Wolf et Móczár, 1972**

MATERIAL. Kozyr': 11.VII 2012, 1 ♀.

DISTRIBUTION. Russia (*Altaysky krai, Irkutskaya oblast, Zabaikalskii krai, Primorskii krai, Amurskaya oblast), Mongolia (Lelej & Loktionov, 2012).

***Arachnospila (Anoplochares) spissa* (Schiodte, 1837)**

Arachnospila (A.) spissa: Tobias, 1978: 132 (key), ♀ ♂; Lelej, 1995: 240 (key), ♀ ♂.

MATERIAL. Kuray: 15.VII 2010, 1 ♂.

DISTRIBUTION. Russia (European part, Altay, Irkutskaya oblast, Zabaikalskii krai, Yakutia, South Sakhalin, Magadanskaya oblast), Western Europe, Ukraine, Belarus, Kazakhstan, Near East (Lelej & Loktionov, 2012).

Arachnospila (Arachnospila) clericalis (F. Morawitz, 1889)

Arachnospila (A.) clericalis: Lelej & Loktionov, 2011: 4, ♀ ♂.

MATERIAL. Kozyr': 11.VII 2012, 3 ♀, 1 ♂; Onguday: 25.VII 2012, 1 ♀, 2 ♂; Chermal: 28.VII 2012, 1 ♀; Karakol: 24.VII 2012, 2 ♀; Chagan-Uzun: 23.VII 2010, 1 ♀; Kuray: 15.VII 2010, 1 ♀; Tigirek: 05. VIII 2011, 1 ♂.

DISTRIBUTION. Russia (Altaysky krai, Tyva, Irkutskaya oblast, Buryatia, Zabaikalskii krai), Turkey, South-East Kazakhstan, Mongolia, China (Gansu) (Lelej & Loktionov, 2011).

Arachnospila (Arachnospila) fumipennis (Zetterstedt, 1838)

Arachnospila (A.) fumipennis (Zetterstedt, 1838): Lelej & Loktionov, 2011: 7, ♀ ♂.

MATERIAL. Tigirek: 05. VIII 2011, 1 ♂.

DISTRIBUTION. Russia (center and south of European part, *Altay Republic, Khakasia, Primorskii krai, Amurskaya oblast, Khabarovskii krai, Magadanskaya oblast, Kamchatka, North Sakhalin), Western Europe, Ukraine, Kazakhstan, Kyrgyzstan, Tajikistan, Mongolia, Near East, China (Hebei), Japan (Hokkaido), North America (Lelej & Loktionov, 2012).

Arachnospila (Arachnospila) rufa (Haupt, 1927)

MATERIAL. Chermal: 28.VII 2012, 1 ♂.

DISTRIBUTION. Russia (centre and south of European part, Orenburgskaya oblast, *Altay Republic, Tyva), Western Europe, Belarus, Turkey, North and Central Kazakhstan, Kyrgyzstan, Mongolia, Near East (Lelej & Loktionov 2011).

Eoferreola manticata (Pallas, 1771)

MATERIAL. Tigirek: 05.VIII 2011, 1 ♀.

DISTRIBUTION. Russia (south of European part, *Altaysky krai, Primorskii krai), Europe, Kazakhstan, Near East, North Africa (Lelej & Loktionov, 2012).

Episyron albonotatum (Vander Linden, 1827)

MATERIAL. Chermal: 28.VII 2012, 1 ♀, 1 ♂; Tigirek: 05.VII 2012, 1 ♀; 05.VIII 2011, 1 ♂; Chaynaya: 09.VII 2012, 1 ♀; Chagan-Uzun: 23.VII 2010, 1 ♂.

DISTRIBUTION. Russia (European part, *Altay Republic, Irkutskaya oblast, Buryatia, Zabaikalskii krai, North Sakhalin, Khabarovskii krai), Europe, Kazakhstan, Central Asia, Mongolia (Lelej & Loktionov, 2012).

***Episyron candiotum* Wahis, 1966**

MATERIAL. Chermal: 28.VII 2012, 1 ♀.

DISTRIBUTION. Russia (North Caucasus, south of European part, *Altay Republic, Irkutskaya oblast, Primorskii krai, Amurskaya oblast), Europe, Turkey, Kazakhstan, Mongolia, Korea, China (Lelej & Loktionov, 2012).

***Evagetes crassicornis* (Shuckard, 1837)**

MATERIAL. Tigirek: 04.VIII 2011, 1 ♂; Kamlak: 18.VII 2012, 1 ♂.

DISTRIBUTION. Russia (European part, *Altaysky krai, Irkutskaya oblast, Buryatia, Zabaikalskii krai, Yakutia, Amurskaya oblast, Chukotka, South Sakhalin, Magadanskaya oblast), Europe, Georgia, Morocco, Kazakhstan, Kyrgyzstan, Tajikistan, Mongolia, South America (Lelej, Loktionov, 2012).

***Evagetes gibbulus* (Lepelletier de Saint Fargeau, 1845)**

MATERIAL. Kolyvansky: 05.VII 2010, 1 ♀.

DISTRIBUTION. Russia (Caucasus, European part, *Altaysky krai), Europe, Kazakhstan, Kyrgyzstan, Mongolia (Loktionov & Lelej, 2009; Wahis, 2013).

***Evagetes proximus* (Dahlbom, 1845)**

MATERIAL. Chermal: 28.VII 2012, 1 ♂; Onguday: 25.VII 2012, 1 ♂.

DISTRIBUTION. Russia (European part, *Altay Republic, Irkutskaya oblast, Buryatia, Zabaikalskii krai, Primorskii krai, Amurskaya oblast, Khabarovskii krai, Magadanskaya oblast), Western Europe, Ukraine, Belarus, Turkey, Iran, Kazakhstan, Kyrgyzstan, Tajikistan, Mongolia, China (Hebei), Japan (Hokkaido) (Loktionov & Lelej, 2009).

***Evagetes subglaber* (Haupt, 1941)**

MATERIAL. Tigirek: 05.VIII 2011, 1 ♀; Kozyr': 11.VII 2012, 4 ♀; Onguday: 25.VII 2012, 1 ♀.

DISTRIBUTION. Russia (center and south of European part, *Altaysky krai), Western Europe, Ukraine, Mongolia, Kyrgyzstan, Turkmenistan (Loktionov & Lelej, 2009).


***Evagetes trispinosus* (Kohl, 1886)**

Evagetes pectinipes trispinosus: v.d. Smissen, 2003: 158, ♀ ♂.

Evagetes trispinosus: Loktionov & Lelej, 2009: 392, ♀ ♂.

DIAGNOSIS. FEMALE. Anterior border of labrum with median notch (Fig. 1). Tarsal comb with flattened setae. First tarsomere apical spine length second and third tarsomeres length combined or little bit shorter. Body with brown pubescence.

Temples and propleura with brown setae. 11–14.8 mm. MALE. First flagellomere 1.4–1.5 times its width. Hypopygium with one median carina. Genitalia as in Fig. 2. The head and body with long rare setae. Head, scape, pronotum, mesonotum and propodeum with pale brown setae. Propodeum with silver pubescence. 7.6–12.5 mm.



Figs 1, 2. *Evagetes trispinosus* (from: v.d. Smissen, 2003). 1 – clypeus of female, frontal view; 2 – male genitalia, ventral view; 3 – the same, lateral view.

MATERIAL. Tigirek: 05. VIII 2011, 1 ♀.

DISTRIBUTION. *Russia (Altaysky krai), Europe (Bulgaria, Greece, Spain, Italy, Cyprus, Portugal, France, Croatia), Turkey, Kazakhstan, Kyrgyzstan, Near East (Lelej & Loktionov, 2012).

***Parabatozonus lacerticida* (Pallas, 1771)**

Batozonellus lacerticida: Tobias, 1978: 118 (key), ♀ ♂; Lelej, 1995: 253 (key), ♀ ♂.

Parabatozonus lacerticida: Loktionov & Lelej, 2014: 213, ♀ ♂.

MATERIAL. Kolyvansky: 05.VII 2010, 1 ♂.

DISTRIBUTION. Russia (European part, Altaysky krai, Irkutskaya oblast, Buryatia, Zabaikalskii krai, Primorskii krai, Amurskaya oblast, Khabarovskii krai), Europe, Turkey, Kazakhstan, Central Asia, Japan (Hokkaido, Honshu), Republic of Korea, North-eastern China, North Africa (Lelej & Loktionov, 2012).

ACKNOWLEDGEMENTS

The author are indebted to Dr. A.S. Lelej, V.M. Loktionov (Institute of Biology and Soil Science, Vladivostok) for the comprehensive assistance, and Dr. S.A. Belokobylskij (Zoological Institute, St. Petersburg) for the permission to work with pompilid collection. The work was supported by the Russian Found of Basic Research (grant No. 12–04–31932).

REFERENCES

- Bagirov, R.T-o., Burmistrov, M.V., Byvaltsev, A.M., Chernyshev, S.E., Costrov, D.V., Danilov, Yu.N., Dudko, A. Yu., Dudko, R.Yu., Filimonov, R.V., Knyshev, A.A., Kosova, O.V., Krugova, T.M., Kuzmenkin, D.V., Kuznetsova, R.O., Legalov, A.A., Lvovsky, A.L., Namyatova, A.A., Nedoshivina, S.V., Perunov, Yu.E., Reschikov, A.V., Shcherbakov, M.V., Sheikin, S.D., Sinev, S.Yu., Solovarov, V.V., Trilikauskas, L.A., Tshesnokova, S.V., Tyumaseva, Z.I., Udalov, I.A., Ustyuzhanin, P.Ya., Vasilenko, S.V., Vishnevskaya, M.S., Volinkin, A.V. & Yanygina, L.V. 2011. Invertebrates of the Tigirek strict nature reserve. *Proceedings of the Tigirek State Natural Reserve. Barnaul*. Vol. 4. P. 165–226. [In Russian].
- Lelej, A.S. 1995. 64. Fam. Pompilidae – Spider wasps. In: Lelej A.S., Kupianskaya A.N., Kurzenko N.V. & Nemkov P.G. (Eds). *Key to the Insects of the Russian Far East. Vol. 4. Pt 1*. St. Petersburg: Nauka. P. 211–264. [In Russian].
- Lelej, A.S. & Loktionov, V.M. 2011. Review of the nominotypical subgenus of *Arachnospila* Kincaid (Hymenoptera: Pompilidae) of Russia and neighbouring countries with the lectotypification of enigmatic *Pompilus sogdianus* Morawitz and description of new species. *Zootaxa*, 2882: 1–18.
- Lelej, A.S. & Loktionov, V.M. 2012. 59. Fam. Pompilidae (*Psammocharidae*) – Spider wasps. In: Lelej A.S. (Ed.). *Annotated catalogue of the insects of Russian Far East. Volume I. Hymenoptera*. Vladivostok: Dalnauka. P. 407–414. [In Russian].
- Loktionov, V.M. & Lelej, A.S. 2009. A review of the genus *Evagetes* Lepeletier, 1845 (Hymenoptera, Pompilidae) of the Russian fauna. *Euroasian entomological journal*, 8(4): 387–398. [In Russian].
- Loktionov, V.M. & Lelej, A.S. 2011. Review of the subgenus *Ammosphex* Wilcke, 1942 of the genus *Arachnospila* Kincaid, 1900 (Hymenoptera: Pompilidae) of the Russian Far East and East Siberia. *Zootaxa*, 3137: 1–30.
- Loktionov, V.M. & Lelej, A.S. 2012. Taxonomic notes on three species of spider wasps (Hymenoptera, Pompilidae) from Russian Far East. *Far Eastern Entomologist*, 243: 10–14.
- Loktionov, V.M. & Lelej, A.S. 2014. *Spider wasps (Hymenoptera: Pompilidae) of the Russian Far East*. Vladivostok: Dalnauka. 472 p. [In Russian].
- Shlyahtenok, A.S., Lelej, A.S. & Loktionov, V.M. 2012. A review of the genus *Anoplius* Dufour, 1834 (Hymenoptera, Pompilidae) of European fauna. *Euroasian entomological journal*, 11(5): 449–464. [In Russian].
- Tobias, V.I. 1978. Superfamily Pompiloidea. In: Medvedev G.S. (Ed.). *Key to the Insects of the European part of the USSR. Vol. 3. Pt 1*. Leningrad: Nauka. P. 83–147. [In Russian].
- Wahis, R. 2013. Fauna Europaea: Hymenoptera, Pompilidae. *Fauna Europaea version 2.6.2*, <http://www.faunaeur.org> (accessed March 4, 2013).
- Van der Smitten, J. 2003. Revision der europäischen und türkischen Arten der Gattung *Evagetes* Lepeletier 1845 unter Berücksichtigung der Geäderabweichungen. Mit zweisprachigem Schlüssel zur Determination (Hymenoptera: Pompilidae). *Verhandlungen des Vereins für Naturwissenschaftliche Heimatforschung zu Hamburg e. V.*, 42. P. 1–253. [In German].

Correspondence

hppt/ urn:lsid:zoobank.org:pub: 4037E1EA-2177-4970-9426-A92B66D0F95C

A. L. Ozerov¹⁾, M. G. Krivosheina²⁾. THE FIRST RECORD OF THE FAMILY CARNIDAE (DIPTERA) IN THAILAND, WITH THE DESCRIPTION OF A NEW SPECIES. – *Far Eastern Entomologist*. 2014. N 279: 11-12.

1) Zoological Museum, Moscow Lomonosov State University, Bol'shaya Nikitskaya 6, Moscow 125009, Russia. E-mail: ozerov2455@rambler.ru

2) A.N. Severtzov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow 119071, Russia. E-mail: dipteramarina@rambler.ru

Summary. *Meoneura thaica* sp. n. is described from Thailand. Holotype of the new species is deposited in Zoological Museum, Moscow State University. The family Carnidae and the genus *Meoneura* Rondani, 1856 are registered in Thailand for the first time.

Key words: Diptera, Carnidae, *Meoneura*, new species, Thailand.

А. Л. Озеров¹⁾, М. Г. Кривошеина²⁾. Первое указание семейства Carnidae (Diptera) для Таиланда с описанием нового вида // Дальневосточный энтомолог. 2014. N 279. С. 11-12.

Резюме. Из Таиланда описан *Meoneura thaica* sp. n. Голотип нового вида хранится в коллекции Зоологического музея МГУ. Семейство Carnidae и род *Meoneura* Rondani, 1856 впервые отмечены для Таиланда.

INTRODUCTION

The majority of the carnid species belong to the genus *Meoneura* Rondani, 1856 and the species of this genus are mostly Holarctic (Brake, 2009). Oriental fauna is studied poorly and includes now 7 species (Papp, 2013): *Carnus orientalis* Maa, 1968 from Malaysia, *Meoneura vikhrevi* Ozerov, 2011, *Meoneura indica* Papp, 2013, and *Meoneura simplex* Papp, 2013 from India, *Meoneura orientalis* Papp, 2013 from Laos, *Meoneura nepalensis* Papp, 2013 from Nepal, and *Meoneura subinversa* Papp, 2013 from Vietnam.

The description of a new species, *Meoneura thaica* sp. n., from Thailand is given below. The family Carnidae and genus *Meoneura* are registered in Thailand for the first time. The holotype of *M. thaica* sp. n. is deposited in the Zoological Museum of Moscow University, Moscow (ZMUM).

DESCRIPTION OF NEW SPECIES

***Meoneura thaica* Ozerov et Krivosheina, sp. n.**

Figs 1, 2

MATERIAL. Holotype – ♂, Thailand: Kanchanaburi province, Taweechai elephant camp, 14,215° N, 99,225° E, 28.I 2014, coll. by N.E. Vikhrev (in ZMUM).

DESCRIPTION. MALE. Body length 1.1 mm; wing length 0.9 mm. Head, thorax, abdomen and legs black, but tarsi of all legs dark brown. Frontal triangle extended nearly to lunula,

shining. Scutum shining rather than subshining, pleural sclerites subshining. No dorsocentral setae. Scutellum microtomentose. Wing hyaline. Knob of halter yellow, stalk blackish. Abdomen subshining. Epandrium with one pair of epandrial processes only (surstyli). Surstylus simple, straight and wide in lateral view (Fig. 1).



Figs 1–2. *Meoneura thaica* sp. n., holotype male: 1) epandrium, cercus and surstylus, lateral view; 2) epandrium, cerci and surstyli, dorsal view.

DIAGNOSIS. Among Oriental species of *Meoneura*, the new species resembles *Meoneura vikhrevi* Ozerov which also has one pair of epandrial processes only, but its surstylus is hook-like in lateral view (see: Ozerov, 2011: 3, fig. 4). Besides, *M. thaica* is characterized by the absence of dorsocentral setae, and *M. vikhrevi* has one pair of dorsocentral setae.

REFERENCES

- Brake, I. 2009. World Catalog of the Family Carnidae (Diptera, Schizophora). *Myia*, 12: 113–169.
- Papp, L. 2013. New species and records of *Meoneura* Rondani, 1856 from the Old World tropics (Diptera, Carnidae). *Acta Zoologica Academiae Scientiarum Hungaricae*, 59(3): 213–227.
- Ozerov, A.L. 2011. Three new species of the genus *Meoneura* Rondani, 1856 (Diptera, Carnidae) from Asia. *Far Eastern Entomologist*, 221: 1–4.

© **Far Eastern entomologist (Far East. entomol.)** Journal published since October 1994.

Editor-in-Chief: S.Yu. Storozhenko

Editorial Board: A.S. Lelej, N.V. Kurzenko, M.G. Ponomarenko, E.A. Beljaev, V.A. Mutin, E.A. Makarchenko, T.M. Tiunova, P.G. Nemkov, M.Yu. Proshchalykin, S.A. Shabalin

Address: Institute of Biology and Soil Science, Far East Branch of Russian Academy of Sciences, 690022, Vladivostok-22, Russia.

E-mail: entomol@ibss.dvo.ru

web-site: <http://www.biosoil.ru/fee>