


Milichiella lacteipennis: new record for Lampedusa Island (Italy)

Alfio RASPI¹, Silvestro PISCIOTTA², Maurizio SAJEVA²

¹Dipartimento di Coltivazione e Difesa delle Specie Legnose "G. Scaramuzzi", Università di Pisa

²Dipartimento di Scienze Botaniche, Università degli Studi di Palermo, Italy



Abstract

The authors report the first record of *Milichiella lacteipennis* (Loew) (Diptera Milichiidae) in Lampedusa Island (Italy), and give information on its distribution and biology.

Key words: Apocynaceae, insect pollinators.

Introduction

In the course of field investigations carried out during a research project on the pollination ecology of two fly-pollinated Apocynaceae in Lampedusa Island, numerous species of Diptera have been identified as pollinators. Among the pollinators, seven species of Diptera new for Lampedusa have been found (Pisciotta *et al.*, 2008). During further field work several specimens of *Milichiella lacteipennis* (Loew) (Diptera Milichiidae) were captured. In the present note we report the detection of this species in Lampedusa Island, as the first record for Italy. Lampedusa Island (Italy) (35°31'46"N and 35°29'35"N - 12°30'54"E and 12°37'55"E) lies in the Mediterranean Sea and is 20.2 Km² - 11 Km long (East-West) with a maximum broadness of 3.5 Km. It is an emergence of the African Continental platform and is 195 Km from the Sicilian coast and 120 Km from Tunisia (Agnesi and Federico, 1995).

Materials and methods

Fieldwork was carried out from April 2006 to November 2008. Most of the observations were diurnal, with limited nocturnal surveillance. Since the main scope of the work was to investigate the pollination ecology of the two species of Apocynaceae co-occurring in Lampedusa (*Periploca laevigata* subsp. *angustifolia* (Labill.) Markgraf and *Caralluma europaea* (Guss.) N.E.Br., captures were carried out following two techniques: (1) stalking near plants visually searching for arrival of probable pollinators, (2) random captures in the sampling area. Captures were done with an entomological net and specimens were kept separately in plastic test-tube with cork shaving imbued with ethyl acetate until they could be dry prepared for identification. The island was visited monthly to observe insect behaviour and to collect specimens for identification. Specimens have been deposited at the Entomological Collection of the Dipartimento di Coltivazione e Difesa delle Specie Legnose "G. Scaramuzzi", University of Pisa and at the Dipartimento di Scienze Botaniche, University of Palermo.



Figure 1. *M. lacteipennis* (length: 2.4 mm) with pollinaria of *C. europaea* attached to its mouthparts. (In colour at www.bulletinofinsectology.org)

Results and discussion

In November 2008, while monitoring insect activity around *C. europaea* plants, several specimens of *M. lacteipennis* were observed. The flies were actively visiting the flowers with the typical behaviour of pollinators visiting stapeliads flowers as reported by Meve and Liede (1994). Thirty specimens of *M. lacteipennis* were captured, and among them twenty four specimens had pollinaria of *C. europaea* attached to their mouthparts (figure 1).

M. lacteipennis, described by Loew (1866) as *Lobioptera lacteipennis* from Cuba.

Specimens examined: Italy, Lampedusa Island, loc. Isola dei Conigli, 7-8.XI.2008, 30 females, Silvestro Pisciotta legit.

The genus *Milichiella* Giglio-Tos 1895 is characterized by an anepisternum (mesopleuron) bare and notched posterior margin of eye. In the Palearctic Region there are 5 described species (Papp, 1984; Carles-Tolrà, 2001). *M. lacteipennis* is characterized by halteres yellowish white, palpus black, mesonotum with only the 2nd pair of posteriorly-located dorsocentral setae strong.

Thorax and the elongate 5th tergite black shining, wing hyaline (figure 1) with veins R₄₋₅ and M₁ strongly convergent apically. This species resembles *Milichiella iberica* Carles-Tolrà (2001) but in *M. lacteipennis* all the legs (including all tarsi) are black and the mesonotum is shiny black, not slightly microtomentose.

The species is widespread in the Nearctic, Neotropical, Afrotropical and Oriental Regions (Papp, 1984), but in the Palaearctic Region is previously known only from the Canary and Madeira Islands (Becker, 1907, 1908a, 1908b; Frey, 1936), Cape Verde Islands (Dawah and Abdullah, 2007), Morocco (Brake, 2000), Tunisia (Deeming *in litteris*: Sidi Boul Ali, olive grove, 9.V.1995, 1 female, now in the collection of National Museum and Galleries of Wales, Cardiff), Egypt (Steyskal and El-Bialy, 1967), Israel (Brake, 2000), Saudi Arabia (Dawah and Abdullah, 2007), Oman (Deeming, 1998), Afghanistan (Papp, 1979) and Japan (Iwasa, 1999).

This species has been reported on the African continent from Nigeria (Sabrosky, 1980) to Kenya and Zambia (Dawah and Abdullah, 2007), from Seychelles to Madagascar and Reunion (Dawah and Abdullah, 2007) and from Ascension Island and St. Helena, (Dawah and Abdullah, 2007).

The species is widespread throughout the New World from Massachusetts (Johnson, 1925) to California (Malloch, 1913), Alabama (Landau and Gaylor, 1987) and Florida (Johnson, 1913) in the north, and from Venezuela (Galan and Herrera, 2006) to Brazil (Malloch, 1913), Peru (Becker, 1907) and Chile (Malloch, 1934) in the south.

The species has been reported from the Oriental and Australian Regions including: Taiwan (Malloch, 1914), New Guinea and Java (Becker, 1907) and from Australia (Malloch, 1931) and the Chesterfield Islands (Cohic, 1959). It has also been reported from several oceanic islands such as the Hawaiian Islands (Hardy and Delfinado, 1980), Fiji (Bezzi, 1928) and Easter Islands (Campos and Peña, 1973).

Adults of Milichiidae visit flowers, carrion and dung, with some species being commensalists of predatory insects and spiders (Papp, 1984; Brake, 2000; Oosterbroek, 2006). Larvae are saprophagous in organic matter in a variety of niches, including: wet media such as rotting fish, soil, rotting wood, snails, dead insects and pupae, decaying vegetable matter and many types of dung and dry media such as grain and bird nest detritus or ant nests, wood detritus and dry guano (Ferrar, 1987). Raspi (Leto Barone *et al.*, 2002) found many puparia of Milichiidae [probably of *Leptomelopa latipes* (Meigen)] in the sarcophagus of Federico II in Palermo.

Bezzi (1928) reported *M. lacteipennis* reared from heaps of stable manure and rotting cow-pea seeds in Fiji. Bohart and Gressitt (1958) reared this species from dung of various animals (cow, horse, chicken, guinea pig) and from decaying aquatic vegetation. The biology of *Milichiella* is consistent with our report of *M. lacteipennis* visiting flowers and acting as a pollinator of *C. europaea*, whose flowers are sapromyophilic (pollinated by carrion flies). The visual attractants and floral scent composition are important in the attraction of pollinators in sapromyophilous systems, where the odour re-

sembles an oviposition site or a food source for carrion feeders in the orders Coleoptera and Diptera (Dobson, 2006; Jürgens *et al.*, 2006).

It is interesting to note that we found *M. lacteipennis* only in November 2008, despite capturing over 1500 specimens of Diptera in Lampedusa Island starting in April 2006. This species should be an old presence in Lampedusa Island for its proximity to Africa, but apparently *M. lacteipennis* was never reported in Lampedusa (Venturi, 1960; Pisciotto *et al.*, 2008), Linosa or Pantelleria Islands (Raspi, 1995) in the past years.

Acknowledgements

We would like to thank the Director G. Nicolini and the staff of Riserva Naturale Orientata Isola di Lampedusa for assistance and logistic support. We are grateful to Dr. Irina Brake (Natural History Museum, London) and two anonymous reviewers for comments and suggestions to the manuscript. We are grateful also to Paolo Giannotti (Dip. Coltivazione e Difesa delle Specie Legnose "G. Scaramuzzi", Pisa) for valuable assistance in preparing the photographic documentation.

References

- AGNESI V., FEDERICO C., 1995.- Aspetti geografico-fisici e geologici di Pantelleria e delle isole Pelagie (Canale di Sicilia).- *Naturalista siciliano*, 19 (Suppl.): 1-22.
- BECKER TH., 1907.- Die Dipteren-Gruppe Milichinae.- *Annales historico-naturales Musei nationalis hungarici*, 5 (2): 507-550.
- BECKER TH., 1908a.- Dipteren der Kanarischen Inseln.- *Mitteilungen aus dem Zoologischen Museum in Berlin*, 4 (1): 1-180.
- BECKER TH., 1908b.- Dipteren der Insel Madeira.- *Mitteilungen aus dem Zoologischen Museum in Berlin*, 4 (1): 181-206.
- BEZZI M., 1928.- *Diptera Brachycera and Athericera of the Fiji Islands*.- British Museum (Natural History), London, UK.
- BOHART G. E., GRESSIT J. L., 1958.- Filth-inhabiting flies of Guam.- *Bernice Pauahi Bishop Museum Bulletin*, 204: 1-152.
- BRAKE I., 2000.- Phylogenetic systematics of the Milichiidae (Diptera, Schizophora).- *Entomologica Scandinavica*, Suppl. 57: 1-120.
- CAMPOS S. L., PEÑA G. L. E., 1973.- Los insectos de Isla de Pascua.- *Revista Chilena de Entomología*, 7: 217-229.
- CARLES-TOLRÀ M., 2001.- A new species of *Milichiella* Giglio-Tos from Spain (Diptera, Milichiidae).- *Studia Dipterologica*, 8 (2): 521-523.
- COHIC F., 1959.- Report on a visit to the Chesterfield Islands, September 1957.- *Atoll Research Bulletin*, 63: 1-11.
- DAWAH H. A., ABDULLAH M. A., 2007.- New records of some filth flies species (Diptera: Milichiidae) in southwest Saudi Arabia.- *Saudi Journal of Biological Sciences*, 14: 263-270.
- DEEMING J. C., 1998.- Milichidae and Carnidae (Diptera: Cyclorrhapha) from the Arabian Peninsula.- *Fauna of Arabia*, 17: 147-157.
- DOBSON H. E. M., 2006.- Relationship between floral fragrance composition and type of pollinator. pp. 147-198. In: *Biology of Floral Scent* (DUDAREVA N. A., PICHERSKY E., Eds).- CRC Press, Boca Raton, USA.
- FERRAR P., 1987.- *A guide to the breeding habits and immature stages of Diptera Cyclorrhapha*.- Entomograph 8 (LYNEBORG L., Ed.).- E. J. Brill/Scandinavian Science Press, Leiden, Copenhagen.

- FREY R., 1936.- Die Dipterenfauna der Kanarischen Inseln und ihre Probleme.- *Acta Societatis Scientiarum Fennicae Commentationes Biologicae*, 6: 1-236.
- GALAN C., HERRERA F. F., 2006.- Fauna cavernicola de Venezuela: una revisión.- *Boletín de la Sociedad Venezolana de Espeleología*, 40: 39-57, 200.
- GIGLIO-TOS H., 1895.- Mission scientifique de M. Ch. Allaud aux îles Séchelles. Dipteres.- *Annales de la Société entomologique de France*, 64: 353-368.
- HARDY D. E., DELFINADO M. D. 1980.- Diptera Cyclorrhapha III, Series Schizophora, Section Acalyptera, exclusive of family Drosophilidae.- In: *Insects of Hawaii*, 13 (LIEBHERR J. K., Ed.).- University of Hawaii Press, Honolulu, USA.
- IWASA M., 1999.- The Genus *Milichiella* Giglio-Tos (Diptera, Milichiidae) of Japan.- *Medical Entomology and Zoology*, 50: 27-32.
- JOHNSON C. W., 1913.- Insects of Florida. I. Diptera.- *Bulletin of the American Museum of Natural History*, 32: 37-90.
- JOHNSON C. W., 1925.- Fauna of New England. 15. List of the Diptera or two-winged flies.- *Occasional Papers of the Boston Society of Natural History*, 7 (15): 1-326.
- JÜRGENS A., DÖTTERL S., MEVE U., 2006.- The chemical nature of fetid floral odours in stapeliads (Apocynaceae-Asclepiadoideae-Ceropegieae).- *New Phytologist*, 172: 452-468.
- LANDAU G. D., GAYLOR M. J., 1987.- Observations on commensal Diptera (Milichiidae and Chloropidae) associated with spiders in Alabama.- *The Journal of Arachnology*, 15: 272.
- LETO BARONE G., LIOTTA G., RASPI A., 2002.- Indagini entomologiche-Metodi e scelte, pp. 107-113.- In "Il Sarcofago dell'imperatore. Studi, ricerche e indagini sulla tomba di Federico II nella Cattedrale di Palermo".- Regione Siciliana. Assessorato dei Beni Culturali ed Ambientali e della Pubblica Istruzione, Palermo, Italy.
- LOEW H., 1866.- Diptera Americae septentrionalis indigena. Centuria Sexta.- *Berliner Entomologische Zeitschrift*, 9 (1865): 127-186.
- MALLOCH J. R., 1913.- A synopsis of the genera of Agromyzidae, with descriptions of new genera and species.- *Proceedings of the United States National Museum*, 46: 127-154.
- MALLOCH J. R., 1914.- Formosan Agromyzidae.- *Annales historico-naturales Musei nationalis hungarici*, 12: 306-336.
- MALLOCH J. R., 1931.- Notes on Australian Diptera.- *Proceedings of the Linnean Society of New South Wales*, 56: 60-78.
- MALLOCH J. R., 1934.- Acalyptera (concluded), pp. 393-489. In: *Diptera of Patagonia and South Chile, Vol. part VI/f*.- British Museum (Natural History), London, UK.
- MEVE U., LIEDE S., 1994.- Floral biology and pollination in stapeliads - new results and a literature review.- *Plant Systematics and Evolution*, 192: 99-116.
- OOSTERBROEK P., 2006.- *The European Families of the Diptera. Identification, diagnosis, biology*.- KNNV Publishing, Utrecht, The Netherlands.
- PAPP L., 1979.- Milichiidae and Carnidae from Afghanistan.- *Folia entomologica hungarica*, 32:105-109.
- PAPP L., 1984.- Family Milichiidae, pp. 110-118. In: *Catalogue of Palearctic Diptera 10, Chusidae, Chloropidae* (SOOS A., PAPP L., Eds).- Elsevier, Amsterdam, The Netherlands.
- PISCIOTTA S., ZITO P., SAJEVA M., RASPI A., 2008.- New records of Diptera for Lampedusa Island.- *Naturalista siciliano*, 32: 343-349.
- RASPI A., 1995.- Diptera.- *Naturalista siciliano*, 29 (suppl.): 667-674.
- SABROSKY C. W., 1980.- 75. Family Milichiidae, pp. 686-689. In: *Catalogue of the Afrotropical Region* (CROSSKEY R. W., Ed.).- British Museum (Natural History), London, UK.
- STEYSKAL G. C., EL-BIALY S., 1967.- A list of Egyptian Diptera with a bibliography and key to families.- *Technical Bulletin 3*. Ministry of Agriculture, Dokki Cairo, UAR.
- VENTURI F., 1960.- Diptera.- In: ZAVATTARI E., 1960.- Biogeografia delle isole Pelagie.- *Rendiconti. Accademia Nazionale dei XL*, Roma, ser. IV, 11: 342-349.

Authors' addresses: Alfio RASPI (corresponding author: araspi@agr.unipi.it), Dipartimento di Coltivazione e Difesa delle Specie Legnose "G. Scaramuzzi", sezione Entomologia Agraria, Università di Pisa, via San Michele degli Scalzi 2, 56124 Pisa, Italy; Silvestro PISCIOTTA, Maurizio SAJEVA, Dipartimento di Scienze Botaniche, Università degli Studi di Palermo, via Archirafi 38, 90123 Palermo, Italy.

Received March 4, 2009. Accepted May 11, 2009.